

PARTICIPATORY EPIDEMIOLOGY TO ASSESS SARCOPTIC MANGE IN SEROW OF TAIWAN

Author(s): Chen-Chih Chen, Kurtis Jai-Chyi Pei, Yu-Ching Lai, and Jack A. Mortenson

Source: Journal of Wildlife Diseases, 48(4):869-875.

Published By: Wildlife Disease Association

DOI: <http://dx.doi.org/10.7589/2011-07-189>

URL: <http://www.bioone.org/doi/full/10.7589/2011-07-189>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

PARTICIPATORY EPIDEMIOLOGY TO ASSESS SARCOPTIC MANGE IN SEROW OF TAIWAN

Chen-Chih Chen,¹ Kurtis Jai-Chyi Pei,^{1,4} Yu-Ching Lai,² and Jack A. Mortenson³

¹ Institute of Wildlife Conservation, National Pingtung University of Science and Technology, 1 Shuefu Rd., Neipu, Pingtung 91201, Taiwan

² Department of Environmental and Hazards-Resistant Design, Huaan University, 1 Huaan Rd., Shihding, New Taipei City 23301, Taiwan

³ United States Department of Agriculture, Veterinary Services, 530 Center St. NE, Suite 335, Salem, Oregon 97301, USA

⁴ Corresponding author (email: kcpei@mail.npust.edu.tw)

ABSTRACT: We used participatory epidemiology (PE) in remote areas to understand the observed distribution and prevalence of infestation by sarcoptic mange mites (*Sarcoptes scabiei*) on wild Formosan serow (*Capricornis swinhoei*) in Taiwan. A semistructured interview protocol was used for 37 interviews during June–December 2008. Serow with skin lesions consistent with sarcoptic mange were reported within a latitudinal range of approximately 24°00'N to 22°40'N on the Central Mountain Range of Taiwan. The observed prevalence was 40–80% in seven of the 19 interview districts. Clinical signs were observed mainly on serow at elevations >1,000 m and most commonly winter (December–February). *Sarcoptes scabiei* has been observed in the infestation area for at least 80 yr. No other wildlife species with similar skin lesions were reported except wild boar. Sarcoptic mange mites on Taiwan serow might prefer a low-temperature environment, but other factors such as physiologic differences among serow populations might be involved in the determination of the northern boundary of the enzootic range. The use of PE to collect enzootic information on sarcoptic mange in wild serow was effective and rapid.

Key words: *Capricornis swinhoei*, participatory epidemiology, *Sarcoptes scabiei*, Formosan serow.

INTRODUCTION

Field observations in the past decade have reported alopecia and skin lesions on Formosan serow (*Capricornis swinhoei*) in Taiwan (Wang, 2005; Wu, 2005). Infestation by Sarcoptic mange mites (*Sarcoptes scabiei*) was confirmed and documented from a serow in southern Taiwan in 2007 (Chen and Pei, 2007).

Serious impact by sarcoptic mange mites on wildlife populations has been reported in cantabrian chamois (*Rupicapra pyrenaica parva*), common wombat (*Vombatus ursinus*), and red foxes (*Vulpes vulpes*). There are 104 mammalian species reported with sarcoptic mange (Fernández-Morán et al., 1997; Bornstein et al., 2001). The impact of sarcoptic mange on wildlife populations can be severe, especially when newly introduced (Bornstein et al., 2001; Pence and Ueckermann, 2002). Among *Capricornis* species, the Japanese serow (*Capricornis crispus*) is the only other serow species known to be affected by the parasite and

exhibit clinical signs. The first case on Kyushu Island, Japan was found in 1995, and another eight suspected cases were reported in 1995 to 1996 on Kyushu Island (M. Baba, pers. comm.). Two additional cases were discovered in Saitama Prefecture, central Honshu Island, in 1996 and 1997 (Takahashi et al., 2001).

Serow are endemic to Taiwan and widely distributed in the mountainous regions of the island (McCullough, 1974; Lue, 1987), from lowland rainforests to alpine meadow and tundra. Because of human activities at lower elevations, they occur more frequently above 1,000 m (Chiang and Pei, 2008). Since 1989, the Formosan serow has been listed as a “Precious and Rare Species” under Taiwan’s Wildlife Conservation Act, and hunting is prohibited. Nevertheless, illegal hunting continues by indigenous people.

Determining the spatial and temporal distribution of sarcoptic mange in serow is valuable to understanding its potential impact on this protected, endemic species.

Skin lesions caused by sarcoptic mange in serow are specific and characterized by alopecia, lichenification, crusting, fissuring, and skin erosions (Chen and Pei, 2007). Because direct assessment of the mite is not practical, we chose to use participatory epidemiology (PE) in aboriginal tribes to collect disease information, including spatial and temporal distribution, historical occurrence, and possible prevalence of sarcoptic mange in wild serow populations.

An assumption of PE is that local community members have rich and detailed knowledge about the animals they observe and harvest. This survey method is rapid, low cost, and especially useful in remote areas (Jost et al., 2007). Furthermore, through the process of interviews, sample collection and multiple independent respondents, information quality can be controlled and validated.

In Taiwan, the majority of indigenous tribes live in remote mountainous areas and have a long tradition of hunting wildlife for cultural and subsistence reasons. Species commonly hunted are wild boar (*Sus scrofa taiwanus*), Reeves' muntjac (*Muntiacus reevesi micrurus*), sambar (*Rusa unicolor swinhoei*), and serow. Forest rangers, many of whom are also tribal members, work for the Forestry Bureau and National Parks of Taiwan. They have many opportunities to observe serow during the course of their work, and they can also provide PE data.

MATERIALS AND METHODS

Interviews were arranged in indigenous communities and at ranger stations within the known distribution range of serow. Hunters and elders in indigenous communities were contacted by the tribal chief and hunting or religious leaders. Rangers for interview were selected by the directors of each forest district office and national park.

We conducted 37 interviews (25 communities belonging to 20 tribal districts and 12 ranger stations) during June–December 2008 (Fig. 1). Interviews were conducted in groups of two to four interviewees simultaneously. Other tribal members often participated and provided additional information during community interviews. Interviews with tribal members fluent




FIGURE 1. Locations of the 2008 participatory epidemiology interviews and interview results identifying the area absent of skin disease and the area where skin disease was observed in the Formosan serow (*Capricornis swinhoei*) on Taiwan. Distribution range of Formosan serow on the Central Mountain Range (large shaded area on left) and Coastal Mountain Range (small shaded area on right) were adopted from Chiang and Pei (2008).

only in their mother languages were facilitated by translators.

A semistructured interview was used to avoid the potential of leading responses and limiting information, which can occur with formally structured interviews (Ferguson and Messier, 1997). The initial question was “Have you ever seen any sick animal in the wild?” No further prompting or guidance was provided, thereby allowing the interviewees to describe their most memorable incidents. If the responses included “skin disease” or “skin lesion” of serow, interviewees were asked to describe the clinical signs and characteristics of lesions that helped to confirm sarcoptic mange.

Local and color topographic maps (scale 1:50,000) showing trails, rivers, mountains, and villages were used to facilitate identification of hunting grounds or mountain trails where serow with skin diseases or lesions were observed or

hunted. Additional information obtained included elevation range, history of observations, prevalence in the enzootic areas, seasonal occurrence, and other species involved. Prevalence of mange in a tribal district was determined by all participants in an interview through group discussion (Jost et al., 2007). If more than one community interview was conducted within a tribal district, the average was used to represent the prevalence of infestation in that district.

If the interviewee did not mention skin diseases or lesions in their first response, pictures of serow with sarcoptic mange were shown to confirm their observations. The spatial distribution of disease occurrence was identified as the area that serow skin disease was observed by interviewees; the area where serow skin disease had not been seen was categorized as the area of disease absence.

RESULTS

Fifty-four elders and hunters were interviewed; 28 from seven districts stated they had observed skin diseases or lesions on serow. Thirty-eight forest rangers were interviewed, 19 of whom, from five ranger stations, also observed skin diseases or lesions on serow. The seven tribal districts and five ranger stations overlapped spatially. This area has an approximate latitudinal range between 24°00'N and 22°40'N in the Central Mountain Range of Taiwan (Fig. 1). Twenty-five elevation ranges were provided by interviewees in the enzootic region. Skin diseases or lesions of serow were most commonly (76%) seen between 1,000 m and 2,000 m, and the two lowest records were around 700 and 800 m (Fig. 2).

Affected serow were observed in all seven tribal districts during winter (December–February). One district had observations in autumn (September–November) and one in spring (March–April). Average observed prevalence of skin lesions in serow in the enzootic range was 57% (range 40–80%, SD=15, $n=7$).

Tribal elders and hunters in five districts indicated this phenomenon had been seen for more than 80 yr in their hunting grounds. In general, elders and hunters had much more personal experience in




FIGURE 2. Frequency distribution of elevations for 25 locations in Taiwan, where Formosan serow (*Capricornis swinhoei*) with skin disease were observed by interviewees.

handling wildlife than forest rangers. Interviewees from three districts reported similar skin diseases or lesions on wild boar but observed rarely. Sarcoptic mange on wild boar in Taiwan has not been confirmed by appropriate sample collection and laboratory diagnosis. There are no other sympatric ruminates (i.e., muntjac and sambar), or carnivores with similar reported skin lesions.

DISCUSSION

Although PE is developed and used originally for rapid assessments of domestic animal health threats (e.g., Mariner and Roeder, 2003; Jost et al., 2007; Azhar et al. 2010), we demonstrated its usefulness in collecting disease information for a wildlife species. The disease characteristics (lesions) described by the participants could be validated through the process of triangulation by comparing with modern veterinary knowledge of sarcoptic mange and sample collection (Jost et al., 2007).

Nevertheless, biases are expected. The prevalence reported in our study might be an underestimate due to the existence of asymptomatic (in early or recovery stage) cases. Conversely, because areas of alopecia reflect the hunters' spotlights and affected serow usually prefer open areas and move less, serow with skin lesions might be more

easily detected than uninfected individuals, resulting in overestimation. A similar behavior change was observed in common wombats (Hartley and English, 2005). Thus, direct estimation of prevalence and the mortality would be valuable.

According to our interviews, the occurrence of sarcoptic mange infestation on serow was observed in a defined area of the Central Mountain Range. Using the north and south latitudinal boundaries and the knowledge that the majority occurred above 1,000 m, we created a projected distribution map for sarcoptic mange infestation on Taiwan serow (Fig. 3). Skin diseases or lesions on serow outside of this region were rarely or never reported.

It is not clear what factors contribute most to the observed spatial and seasonal patterns we discovered. Poor health status, increased population density, and host susceptibility increase the severity and prevalence of infestation (Arlian, 1989; Fernández-Morán et al., 1997; Skerratt, 2001). Although the low population density at elevations <1,000 m (Chiang and Pei, 2008) might significantly reduce prevalence and frequency of observation, climatic factors such as temperature and humidity may also play a role. All life stages of *S. scabiei* var. *canis* live longer off the host at lower temperatures (10–15 C) and higher humidity (>75%), especially adult females and nymphs (Arlian et al., 1989). Winter was the season sarcoptic mange was most frequently reported in this study and in other hosts (Fernández-Morán et al., 1997; Perez et al., 1997; Skerratt, 2001). Although the optimum climatic conditions for *S. scabiei* on serow are not known, the projected range for the mite in this study has an annual mean minimum temperature <12 C (Fig. 3). If climatic factors are important, it may explain the absence of observed sarcoptic mange on serow in the Coastal Mountain Range and at elevations <700 m in the Central Mountain Range and the lack of southern expansion beyond the known historic range.




FIGURE 3. Projected enzootic range of sarcoptic mange in Formosan serow (*Capricornis swinhoei*) on Taiwan compared to the 12 C annual mean minimum isotherm.

Global mean temperature is increasing (IPCC, 2007); this trend was also observed in Taiwan (Hsu and Chen, 2002). In the past 100 yr, the temperature has increased 1.0–1.4 C in Taiwan and increasing temperature is projected to continue under several climate change scenarios (Hsu and Chen, 2002). Warming temperatures can expand (or shift) the spatial and temporal distribution of pathogens or increase the intensity of disease occurrences (Lafferty, 2009). However, warming temperatures might also decrease the occurrence or have limited effects on a subset of pathogens. Those include pathogens that occur in high elevations or latitudes or that prefer cooler temperatures and directly transmitted pathogens (Harvell et al., 2002; Rohr et al., 2011). Therefore, climate warming could decrease the intensity of sarcoptic mange in serow populations and its spatial and

temporal distribution. Nevertheless, the influence of climate change on other factors, such as habitat loss, host abundance, and host susceptibility, might indirectly affect disease occurrence (Harvell et al., 2002; Lafferty, 2009).

At the northern boundary of the projected range, however, the climate is favorable and there is no evidence to suggest serow population density or general health are factors limiting the northern distribution (Chiang and Pei, 2008; Pei, pers. obs.). It is also unlikely that *S. scabiei* is dispersing northward given the long history of observations in the enzootic range.

A recent phylogenetic study of serow in Taiwan identified two genetically distinct groups with a geographic boundary at the Zhuoshui and Mugua rivers (Horng et al., 2003). These rivers are approximately at the northern boundary of the projected sarcoptic mange enzootic range (Fig. 3). It is possible that the geographic barrier responsible for the distinct serow populations also acts as a barrier to mite dispersal. A second more likely explanation is that serow susceptibility to mites is lower in the northern population. A similar spatial phenomenon was reported in coyote populations in Texas, USA, where two populations had different susceptibility to sarcoptic mange (Pence and Windberg, 1994).

Wild boar was the only other wildlife species observed with skin lesions. However, there were very few observations and no direct evidence to confirm sarcoptic mange. Other sympatric wildlife species have not been found with similar skin lesions in Taiwan. Sarcoptic mange is considered to be caused by a single species of mite that diverged into many morphologically indistinguishable strains on different host species. These strains usually show a high degree of host specificity and little cross infectivity (Bornstein et al., 2001; Pence and Ueckermann, 2002). This has also been demonstrated in the laboratory (Samuel, 1981; Arlian et al., 1988; Arlian et al., 1996). Infestation by *S. scabiei* on serow in Taiwan

might be host specific. Further study of phylogenetic relationships and potential cross-species infestation by *S. scabiei* strains isolated from different wild and domestic species in Taiwan is needed (Walton et al., 2004; Soglia et al., 2007).

As a point of comparison, enzootic information on *S. scabiei* reported in this study is quite different from that in Japan. Many wildlife species in Japan have been identified with sarcoptic mange (Wildlife Management Office, Inc., 1998), including raccoon dog (*Nyctereutes procyonoides*), red fox, wild boar, Japanese badger (*Meles anakuma*), sika deer (*Cervus nippon*), masked palm civet (*Paguma larvata*), and raccoon (*Procyon lotor*). However, only a few cases of sarcoptic mange (Takahashi et al., 2001) have been confirmed in Japanese serow. The solitary, territorial, and predominantly monogamous behaviors of Japanese serow (Kishimoto and Kawamichi, 1996) may limit transmission of the mite.

ACKNOWLEDGMENTS

We thank Yu-Ren Liang for his help in interviewing. Funding was provided by the Taiwan Forest Bureau, Council of Agriculture (Grant 97 Agriculture Technology-11.1.4-Management-e1).

LITERATURE CITED

- ARLIAN, L. G. 1989. Biology, host relations, and epidemiology of *Sarcoptes scabiei*. Annual Review of Entomology 34: 139–159.
- , D. L. V. MOHER, AND D. CORDOVA. 1988. Host specificity of *S. scabiei* var. *canis* (Acari: Sarcoptidae) and the role of host odor. Journal of Medical Entomology 25: 52–56.
- , ———, AND M. J. POLE. 1989. Survival of adults and developmental stages of *Sarcoptes scabiei* var. *canis* when off the host. Experimental and Applied Acarology 6: 181–187.
- , M. S. MORGAN, AND J. J. ARENDS. 1996. Immunologic cross-reactivity among various strains of *Sarcoptes scabiei*. Journal of Parasitology 82: 66–72.
- AZHAR, M., A. S. LUBIS, E. S. SIREGAR, R. G. ALDERS, E. BRUM, J. McGRANE, I. MORGAN, AND P. ROEDER. 2010. Participatory disease surveillance and response in Indonesia: Strengthening veterinary services and empowering communities to

- prevent and control highly pathogenic avian influenza. *Avian Diseases* 54: 749–753.
- BORNSTEIN, S., T. MÖRNER, AND W. M. SAMUEL. 2001. *Sarcoptes scabiei* and sarcoptic mange. In *Parasitic diseases of wild mammals*, 2nd Edition, W. M. Samuel, M. J. Pybus and A. A. Kocan (eds.). Iowa State University Press, Ames, Iowa, pp. 107–119.
- CHEN, C. C., AND K. J. C. PEI. 2007. Case report: The first report of mange infestation in wild Formosan serows (*Capricornis swinhoei*) in Taiwan. *Taiwan Veterinary Journal* 33: 181–185. [In Chinese with English Abstract].
- CHIANG, P. J., AND K. J.-C. PEI. 2008. *Capricornis swinhoei*. In *International Union for Conservation of Nature (IUCN) 2010. IUCN Red List of Threatened Species, Version 2010.4*, www.iucnredlist.org/details/3810/0. Accessed July 2012.
- FERGUSON, M., AND F. MESSIER. 1997. Collection and analysis of traditional ecological knowledge about a population of Arctic tundra caribou. *Arctic* 50: 17–28.
- FERNÁNDEZ-MORÁN, J., S. GÓMEZ, F. BALLESTEROS, P. QUIRÓS, J. L. BENITO, C. FELIU, AND J. M. NIETO. 1997. Epizootiology of sarcoptic mange in a population of cantabrian chamois (*Rupicapra pyrenaica parva*) in Northwestern Spain. *Veterinary Parasitology* 73: 163–171.
- HARTLEY, M., AND A. ENGLISH. 2005. *Sarcoptes scabiei* var. *wombati* infection in the common wombat (*Vombatus ursinus*). *European Journal of Wildlife Research* 51: 117–121.
- HARVELL, C. D., C. E. MITCHELL, J. R. WARD, S. ALTIZER, A. P. DOBSON, R. S. OSTFELD, AND W. M. SAMUEL. 2002. Climate warming and disease risks for terrestrial and marine biota. *Science* 296: 2158.
- HORNG, D. C., H. W. HUANG, Y. C. LIANG, AND B. R. OU. 2003. Two distinct phylogenetic groups of Formosan serow (*Naemohedus swinhoei* Grey) populations in Taiwan: Based on mitochondrial D-loop region sequences. *Endemic Species Research* 5: 15–25. [In Chinese with English Abstract].
- HSU, H. H., AND C. T. CHEN. 2002. Observed and projected climate change in Taiwan. *Meteorology and Atmospheric Physics* 79: 87–104.
- INTERGOVERNMENTAL PANEL ON CLIMATE CHANGE (IPCC). 2007. Summary for Policymakers. In *Climate change 2007: The physical science basis. Contribution of working group I to the fourth assessment report of the Intergovernmental Panel on Climate Change*, S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor, and H. L. Miller (eds.). Cambridge University Press, Cambridge, United Kingdom, pp. 1–18.
- JOST, C. C., J. C. MARINER, P. L. ROEDER, E. SAWITRI, AND G. J. MACGREGOR-SKINNER. 2007. Participatory epidemiology in disease surveillance and research. *Revue Scientifique et Technique-Office International des Epizooties* 26: 537–549.
- KISHIMOTO, R., AND T. KAWAMICHI. 1996. Territoriality and monogamous pairs in a solitary ungulate, the Japanese serow, *Capricornis crispus*. *Animal Behaviour* 52: 673–682.
- LAFFERTY, K. D. 2009. The ecology of climate change and infectious diseases. *Ecology* 90: 888–900.
- LUE, K. Y. 1987. A preliminary study on the ecology of Formosan serow *Capricornis crispus swinhoei*. In *The biology and management of Capricornis and related mountain antelopes*. H. Soma (ed.). Croom Helm, London, UK, pp. 125–133.
- MARINER, J. C., AND P. L. ROEDER. 2003. Use of participatory epidemiology in studies of the persistence of lineage 2 rinderpest virus in East Africa. *Veterinary Record* 152: 641–647.
- MCCULLOUGH, D. R. 1974. Status of larger mammals in Taiwan. *Tourism Bureau, Taipei, Taiwan*, 35 pp.
- PENCE, D. B., AND E. UECKERMANN. 2002. Sarcoptic mange in wildlife. *Revue Scientifique et Technique-Office International des Epizooties* 21: 385–398.
- , AND L. A. WINDBERG. 1994. Impact of a sarcoptic mange epizootic on a coyote population. *The Journal of Wildlife Management* 58: 624–633.
- PÉREZ, J. M., I. RUIZ, J. E. GRANADOS, R. C. SORIGUER, AND P. FANDOS. 1997. The dynamics of sarcoptic mange in the ibex population of Sierra Nevada in Spain. Influence of climatic factors. *Journal of Wildlife Research* 2: 86–89.
- ROHR, J. R., A. P. DOBSON, P. T. J. JOHNSON, A. M. KILPATRICK, S. H. PAULL, T. R. RAFFEL, D. RUIZ-MORENO, AND M. B. THOMAS. 2011. Frontiers in climate change-disease research. *Trends in Ecology and Evolution* 26: 270–277.
- SAMUEL, W. M. 1981. Attempted experimental transfer of sarcoptic mange (*Sarcoptes scabiei*, Acarina: Sarcoptidae) among red fox, coyote, wolf and dog. *Journal of Wildlife Diseases* 17: 343–347.
- SKERRATT, L. F. 2001. Sarcoptic mange in the common wombat, *Vombatus ursinus* (Shaw, 1800). PhD Thesis, University of Melbourne, Melbourne, Australia, 284 pp.
- SOGLIA, D., R. RASERO, L. ROSSI, S. SARTORE, P. SACCHI, AND S. MAIONE. 2007. Microsatellites as markers for comparison among different populations of *Sarcoptes scabiei*. *International Journal of Animal Sciences* 7: 214–216.
- TAKAHASHI, M., S. NOGAMI, H. MISUMI, M. MATSUMOTO, M. TAKAHAMA, AND K. UCHIKAWA. 2001. Mixed infestation of sarcoptic and chorioptic mange mites in Japanese serow, *Capricornis crispus* Temminck, 1845 in Japan, with a description of *Chorioptes japonensis* sp. nov. (Acari: Psorop-

- tidia). *Medical Entomology and Zoology* 52: 297–306.
- WALTON, S. F., A. DOUGALL, S. PIZZUTTO, D. HOLT, D. TAPLIN, L. G. ARLIAN, M. MORGAN, B. J. CURRIE, AND D. J. KEMP. 2004. Genetic epidemiology of *Sarcoptes scabiei* (Acari: Sarcoptidae) in northern Australia. *International Journal for Parasitology* 34: 839–849.
- WANG, Y. 2005. Wildlife monitoring and aborigine hunting at Danta major wildlife habitat. Forest Bureau, Council of Agriculture, Executive Yuan, Taipei, Taiwan, 78 pp. Chinese.
- WILDLIFE MANAGEMENT OFFICE, INC. 1998. An urgent report of the epidemiological survey on sarcoptic mange of Japanese mammals. Ministry of the Environment, Tokyo, Japan, 64 pp. Japanese.
- WU, H. Y. 2005. The study on the Asian black bear and the Artiodactyla community at the eastern area of Yushan National Park. Yushan National Park Headquarters, Department of Interior, Taipei, Taiwan, 59 pp. Chinese.

Submitted for publication 6 July 2011.

Accepted 25 April 2012.